

Table 14. Disease Resistance of Strawberry Cultivars Commonly Grown in the Midwest.

Cultivar	Red Stele	Verticillium Wilt	Leaf Spot	Leaf Scorch	Powdery Mildew
Junebearing					
Allstar	VR ¹	R	R	R	R
Annapolis	S	I	S	S	—
Blomidon	—	—	—	—	—
Canoga	I	I	R	R	—
Cardinal	S	S	R	R	R
Catskill	S	VR	S	R	R
Cavendish	R	I	—	—	S
Delite	R ²	R	R	S-R	S
Earliglow	R ²	T-R	S-R	R	I
Guardian	R ²	T-R	S-R	R	S-R
Honeoye	S	S	R	R	—
Jewel	S	S	R	R	—
Kent	—	—	—	—	—
Lateglow	R	R	T	T	T
Lester	R	R	R	R	R
Midway	R ²	S-I	S	S	I
Pocahontas	S	S	S-R	S-I	R
Raritan	S	S	S	S	I
Redchief	R ²	R	S-R	R	S-R
Scott	R	I-R	S-R	R	R
Seneca	S	S	—	—	—
Sparkle	S-R	S	S	S-I	R
Sunrise	R ²	R	VS	R	R
Surecrop	R ²	VR	S-R	S-R	—
TeMesseeBeauty	S	R	R	S-R	S
Veestar	S	T	T	T	—
Everbearing					
Tribute	VR	T-R	T	T	R
Tristar	R	R	T	T	R

¹ VS=very susceptible; S=susceptible; I=intermediate; T=tolerant; R=resistant; VR=very resistant; —=unknown. Resistant characteristics of the cultivar usually preclude the need for other controls.

²Resistant to several races of the red stele fungus.