

F-0051

# Relative Effectiveness of Herbicides Commonly Used to Control Woody Vegetation in Forest Stands

F-0051-Table 2

Table 2. Relative effectiveness in controlling woody vegetation of selected oil-soluble herbicides applied as a basal spray or cut stump treatment according to label recommendations. <sup>1</sup>			
Species	Triclopyr (e.g., Garlon 4)	Imazapyr (e.g., Chopper, Stalker, & Arsenal)	2,4-D + 2,4-DP (e.g., Patron 170)
Alder	S	S	I-R
Ash, White	S	S	R
Aspen	S	S	S
Balsam Poplar	S	S	S-I
Beech, American	S-I	I-S	S-I
Birch	S	S	I-S
Boxelder	I-S	S	I
Cherry	S-I	S	I-R
Cottonwood	S-I	S	R
Dogwood	S-I	S	I-R
Elderberry	S	S	S-I
Elms	S-I	I-R	S-I
Grapes, Wild	S-I	S	I
Hackberry	I-R	R	S
Hawthorn	S-I	S	I
Honeylocust	S	I-S	I

continued next page

Species	Triclopyr (e.g., Garlon 4)	Imazapyr (e.g., Chopper, Stalker, & Arsenal)	2,4-D + 2,4-DP (e.g., Patron 170)
Honeysuckle	S	S	S
Locust, Black	S	R	I
Maple	S-I	S	I
Maple, Red	S-I	S	R
Maple, Silver	S	S	I
Maple, Sugar	S	S	I
Oaks	I-R	S	S
Osage Orange	R	R	I-R
Persimmon, E.	S	S	S-I
Plum	I-R	S	S-I
Poison Ivy	I-S	S	I
Poplar, Yellow	S-I	S	S
Rose, Multiflora	S-I	S	I
Sassafras	S-I	S	I-R
Sumac	I	S	I-R
Sweetgum	S-I	S-I	R
Sycamore	S	S-I	S-I
Tree of Heaven	S	S	I-R
Trumpetcreeper	I-R	S	R
Virginia Creeper	S	S	R
Willow	S	S	S